

The role of the quark and gluon GPDs in hard vector-meson electroproduction

S.V. Goloskokov ¹

*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,
Dubna 141980, Moscow region, Russia*

P. Kroll ²

Fachbereich Physik, Universität Wuppertal, D-42097 Wuppertal, Germany

Abstract

Electroproduction of light vector mesons is analyzed on the basis of handbag factorization. The required generalized parton distributions are constructed from the CTEQ6 parton distributions with the help of double distributions. The partonic subprocesses are calculated within the modified perturbative approach. The present work extends our previous analysis of the longitudinal cross section to the transverse one and other observables related to both the corresponding amplitudes. Our results are compared to recent experimental findings in detail.

1 Introduction

In a previous work [1] we analyzed electroproduction of light vector mesons ($V = \rho^0, \phi, \omega$) at HERA kinematics within the handbag factorization scheme which is based on generalized parton distributions (GPDs) and hard partonic subprocesses. The latter are calculated within the modified perturbative approach [2] in which the quark transverse momenta are retained. The emission and reabsorption of partons from the proton is still treated in collinear approximation. In the kinematical region accessible to the HERA experiments that is characterized by Bjorken- x (x_{Bj}) of the order 10^{-3} , it is not unjustified to restrict oneself to the gluonic subprocess $\gamma^* g \rightarrow V g$ and the associated gluonic GPD H^g . In a recent paper [3] we extended that analysis to larger values of x_{Bj} ($\lesssim 0.2$) such as accessible to the

¹Email: goloskkv@theor.jinr.ru

²Email: kroll@physik.uni-wuppertal.de




Figure 1: Typical lowest order Feynman graphs for the $\gamma^*g \rightarrow Vg$ (left) and $\gamma^*q \rightarrow Vq$ (right) subprocesses of meson electroproduction.

HERMES experiment, but restricting ourselves to the analysis of the least model-dependent amplitude, namely the one for transitions from longitudinal polarized virtual photons to vector mesons polarized in the same manner, $\gamma_L^*p \rightarrow V_L p$. This analysis necessitates the inclusion of the quark subprocesses $\gamma^*q \rightarrow Vq$ (see Fig. 1) and the associated GPDs for sea and valence quarks.

Here, in this work we are going to complete the analysis of vector-meson electroproduction by studying the amplitude for transversely polarized photons and mesons, $\gamma_T^*p \rightarrow V_T p$. The basic idea of modeling this amplitude has been already described in Ref. [1] for the gluonic contribution. The extension to the quark contribution is straightforward. The crucial point at issue with the transverse amplitude is that the quark transverse momenta which are retained in the modified perturbative approach in order to suppress configurations with large transverse separations of the quark and antiquark forming the meson, regularize the infrared singularities occurring in the transverse subprocess amplitude in collinear approximation [4, 5] at the same time. These special configurations seem to be responsible for the excess of the leading-twist contribution to the longitudinal cross section over experiment. Indeed, taking into account the quark transverse momenta fair agreement between theory and experiment is achieved [3]. We are going to apply this modified handbag approach to vector-meson electroproduction for energies, W , in the photon-proton center of mass system (c.m.s.) between about 5 and 170 GeV and photon virtualities, Q^2 , between about 3 and 100 GeV² while x_{Bj} is less than $\lesssim 0.2$. We will compare our results in detail with recent data from HERA, COMPASS and HERMES.

The GPDs H for quarks and gluons which dominate the process of interest for unpolarized protons at small x_{Bj} , are constructed from the CTEQ6 parton distribution functions (PDFs) [6] through double distributions [7, 8]. Applying the same model to the GPDs \tilde{H} we are also going to estimate the so-called unnatural parity amplitudes and to study their implications on spin density matrix elements (SDME) and the double spin asymmetry A_{LL} describing the correlation of the helicities of the beam and target particles.

In Ref. [9] the electromagnetic form factors of the nucleon have been used to extract the zero-skewness GPDs H , \tilde{H} and E for valence quarks. The forward limit of E_{val}^a , the

analog of the PDFs, determined in Ref. [9], can be utilized for the construction of E_{val}^a at non-zero skewness with the double distribution model. With these GPDs at disposal we will also estimate the SDMEs measurable with a transversely polarized target as well as the transverse spin asymmetry A_{UT} .

The plan of the paper is as follows: In Sect. 2 we will sketch the modified handbag approach. In Sect. 3 the double distribution model for the GPD H is described. The results on cross sections and SDMEs obtained within the handbag approach are presented in Sect. 4 and 5, respectively. Sect. 6 is devoted to an estimate of the role of the GPDs \widetilde{H} and Sect. 7 to that of the GPDs E . We will conclude this paper by a summary (Sect. 8).

2 The handbag approach

We are interested in vector-meson electroproduction in a kinematical region characterized by large Q^2 and large W but small $x_{\text{Bj}} (\lesssim 0.2)$ and small invariant momentum transfer $-t$. In the handbag approach, the amplitudes for the process $\gamma^* p \rightarrow V p$ which can be extracted from vector-meson electroproduction applying the one-photon approximation, factorize into partonic subprocesses (see Fig. 1) and GPDs comprising the soft, non-perturbative QCD. At large Q^2 the amplitude for $\gamma_L^* p \rightarrow V_L p$ dominates and factorization has been shown to hold for it rigorously [10, 11]. The amplitudes for other photon-meson transitions are suppressed by inverse powers of Q . Besides the longitudinal amplitude we will consider only the transverse one, $\gamma_T^* p \rightarrow V_T p$ in this work which is the most important one of the suppressed amplitudes at small $-t$. Proton helicity flip is suppressed by $\sqrt{-t}/2m$ (m being the proton's mass) and can be neglected in calculations of cross sections and SDMEs obtained with unpolarized protons. In Sect. 7 we will however estimate the size of the proton helicity-flip amplitudes explicitly. It will turn out that these amplitudes are indeed small.

In the region of small x_{Bj} the dominant contributions are provided by the GPD H . To the proton helicity non-flip amplitude the GPDs contribute in the combination

$$H - \frac{\xi^2}{1 - \xi^2} E \quad (1)$$

where the skewness ξ is related to x_{Bj} by

$$\xi \simeq \frac{x_{\text{Bj}}}{2 - x_{\text{Bj}}} [1 + m_V^2/Q^2]. \quad (2)$$

Here, m_V denotes the mass of the vector meson. The GPD E can therefore safely be ignored since it is not expected that it is much larger than the GPD H , see Sect. 7 where we will take up this issue again. The GPD \widetilde{H} only contributes to the transverse amplitude and can also be neglected in calculations of the cross sections. In Sect. 6 we will return to this problem and estimate the size of its contribution. Since there is no parton helicity flip in the partonic subprocesses to the accuracy we are calculating them, the parton helicity-flip GPDs [12] do not occur.

The contributions from H to the $\gamma^*p \rightarrow Vp$ amplitudes read ($i = g, q, x_g = 0, x_q = -1$)

$$\mathcal{M}_{\mu+, \mu+}^{Ni}(V) = \frac{e}{2} \sum_a e_a \mathcal{C}_V^a \int_{\bar{x}_i}^1 d\bar{x} \sum_{\lambda} \mathcal{H}_{\mu\lambda, \mu\lambda}^{Vi}(\bar{x}, \xi, Q^2, t=0) H^i(\bar{x}, \xi, t). \quad (3)$$

The first sum runs over the quark flavors a and e_a denotes the quark charges in units of the positron charge e . The non-zero flavor weight factors, \mathcal{C}_V^a , read

$$\mathcal{C}_\rho^u = -\mathcal{C}_\rho^d = \mathcal{C}_\omega^u = \mathcal{C}_\omega^d = 1/\sqrt{2}, \quad \mathcal{C}_\phi^s = 1. \quad (4)$$

The explicit helicities in (3) refer to the proton while μ is the helicity of the photon and meson and λ that of the partons participating in the subprocess. Only the t dependence of the GPDs is taken into account in the amplitudes (3). That of the subprocess amplitudes \mathcal{H} provides corrections of order t/Q^2 which we neglect throughout this paper. In contrast to the subprocess amplitudes the t dependence of the GPDs is scaled by a soft parameter, actually by the slope of the diffraction peak.

There is a minimal value of $-t$ allowed in the process of interest

$$t_{\min} = -4m^2 \frac{\xi^2}{1 - \xi^2}. \quad (5)$$

As other effects of order ξ^2 (e.g. the GPD E) the t_{\min} effect is neglected. We note in passing that our helicities are light-cone ones which naturally occur in the handbag approach. The differences to the usual c.m.s. helicities are of order $m\sqrt{-t}/W^2$ [12] and can be ignored in the kinematical region of interest in this work.

The full amplitude is given by a superposition of the gluon and quark contributions

$$\mathcal{M}^N = \mathcal{M}^{Ng} + \mathcal{M}^{Nq} \quad (6)$$

and it is normalized such that the partial cross sections for $\gamma^*p \rightarrow Vp$ read (Λ is the usual Mandelstam function)

$$\frac{d\sigma_{L(T)}}{dt} = \frac{1}{16\pi(W^2 - m^2)\sqrt{\Lambda(W^2, -Q^2, m^2)}} |\mathcal{M}_{0(+) +, 0(+) +}^N|^2, \quad (7)$$

which holds with regard to the above-mentioned simplifications. The cross sections integrated over t are denoted by σ_L and σ_T . The full (unseparated) cross section for $\gamma^*p \rightarrow Vp$ is

$$\sigma = \sigma_T + \varepsilon\sigma_L, \quad (8)$$

in which ε is the ratio of longitudinal to transverse photon fluxes. The power corrections of kinematical origin given in Eq. (2) and in the phase space factor (7) are taken into account by us. With the exception of these kinematical effects hadron masses are omitted otherwise. In the expression (7) for the cross section the symmetry relation

$$\mathcal{M}_{-\mu\nu', -\mu\nu}^{Ni} = \mathcal{M}_{\mu\nu', \mu\nu}^{Ni} \quad (9)$$

has been used which is an obvious consequence of the definition (3) and parity conservation. This symmetry relation coincides with the one that holds for natural parity exchanges; we therefore mark this amplitude by a superscript N . Since the contributions from \widetilde{H} to the amplitudes obey the relation

$$\mathcal{M}_{-\mu\nu', -\mu\nu}^{Ui} = -\mathcal{M}_{\mu\nu', \mu\nu}^{Ui}, \quad (10)$$

there are no interference terms between \mathcal{M}^{Ni} and \mathcal{M}^{Ui} in the observables for unpolarized electroproduction of vector mesons as can easily be shown. The relation (10) is also obtained for the exchange of a particle with unnatural parity. In analogy to the contributions from H the amplitudes related to \widetilde{H} are marked by a superscript U . The contribution $|\mathcal{M}^U|^2$ is neglected in (7).

Let us now turn to the discussion of the subprocess amplitudes. As is well-known, for the kinematics accessible to current experiments, the handbag amplitude evaluated in collinear approximation, overestimates the longitudinal cross section although with the tendency of approaching experiment with increasing Q^2 [1, 13, 14]. One may wonder whether higher order perturbative QCD corrections to the subprocesses may cure that deficiency. However, this does not seem to be the case. NLO corrections [15, 16] are very large due to BFKL-type logarithms $\sim \ln 1/\xi$ and cancel to a large extent the LO term at low Q^2 and low x_{Bj} . A recent attempt [17] to resum higher orders with methods known from deep inelastic lepton-nucleon scattering [18] seems to indicate that the sum of all higher order corrections to the LO term is not large. In view of this unsettled situation it seems to be reasonable to proceed along the lines advocated in Refs. [1, 3] by simply using the LO result and add power corrections, especially since such corrections are anyway needed in order to account for the large transverse cross section σ_T . Once the higher order perturbative corrections are better understood within the modified perturbative approach one may add them to the LO results. As long as they are of reasonable magnitude there is no principal difficulty in this. It may merely be necessary to readjust the GPDs and the meson wave functions appropriately.

As in our papers [1, 3] we will model the required power corrections by employing the modified perturbative approach [2] in the calculation of the subprocesses. In this approach the transverse momenta of the quark and antiquark, \mathbf{k}_\perp (defined with respect to the meson's momentum), entering the meson are kept. In contrast to the situation at the mesonic vertex, the partons entering the subprocess, are viewed as being emitted and reabsorbed by the proton collinearly. This scenario is supported by the fact that the GPDs describe the full proton and their \mathbf{k}_\perp dependence therefore reflects the proton's charge radius ($\langle \mathbf{k}_\perp^2 \rangle^{1/2} \simeq 200$ MeV) while the meson is generated through its compact valence Fock state with a r.m.s. \mathbf{k}_\perp of about 500 MeV [19, 20]. Instead of a meson's distribution amplitude allowance is to be made for a meson light-cone wavefunction $\Psi_{VL}(\tau, k_\perp)$ in the modified perturbative approach [19, 20]. Here τ is the fraction of the light-cone plus component of the meson's momentum, q' , the quark carries; the antiquark carries the fraction $\bar{\tau} = 1 - \tau$. Quark transverse momenta are accompanied by gluon radiation. In Ref. [2] the gluon radiation has been calculated in the form of a Sudakov factor $\exp[-S(\tau, \mathbf{b}, Q^2)]$

to next-to-leading-log approximation using resummation techniques and having recourse to the renormalization group. The quark-antiquark separation, \mathbf{b} , in configuration space acts as an infrared cut-off parameter. Radiative gluons with wave lengths between the infrared cut-off and a lower limit (related to the hard scale Q^2) yield suppression, softer gluons are part of the meson wavefunction while harder ones are an explicit part of the subprocess amplitude. Congruously, the factorization scale is given by the quark-antiquark separation, $\mu_F = 1/b$, in the modified perturbative approach. For more details we refer to Refs. [2, 3].

Since the resummation of the logarithms involved in the Sudakov factor can only efficiently be performed in the impact parameter space [2] we have to Fourier transform the lowest-order subprocess amplitudes to that space and to multiply them with the Sudakov factor there. This leads to

$$\mathcal{H}_{\mu\lambda,\mu\lambda}^{Vi} = \int d\tau d^2b \hat{\Psi}_{VL}(\tau, -\mathbf{b}) \hat{\mathcal{F}}_{\mu\lambda,\mu\lambda}^i(\bar{x}, \xi, \tau, Q^2, \mathbf{b}) \alpha_s(\mu_R) \exp[-S(\tau, \mathbf{b}, Q^2)]. \quad (11)$$

The two-dimensional Fourier transformation between the canonically conjugated \mathbf{b} and \mathbf{k}_\perp spaces is defined by

$$\hat{f}(\mathbf{b}) = \frac{1}{(2\pi)^2} \int d^2\mathbf{k}_\perp \exp[-i\mathbf{k}_\perp \cdot \mathbf{b}] f(\mathbf{k}_\perp). \quad (12)$$

The renormalization scale μ_R is taken to be the largest mass scale appearing in the hard scattering amplitude, i.e. $\mu_R = \max(\tau Q, \bar{\tau} Q, 1/b)$. Since the bulk of the handbag contribution to the amplitudes is accumulated in regions where μ_R is smaller than 3 GeV we have to deal with three active flavors. A value of 220 MeV for Λ_{QCD} is used in the Sudakov factor and in the evaluation of α_s from the one-loop expression.

The hard scattering kernels \mathcal{F}^i or their Fourier transform $\hat{\mathcal{F}}^i$ occurring in Eq. (11), are computed from the pertinent Feynman graphs, see Fig. 1. The result for the gluonic subprocess is discussed in some detail in Ref. [1] and we refrain from repeating the lengthy expressions here. For quarks, on the other hand, the hard scattering kernel for longitudinally polarized photons and mesons reads

$$\mathcal{F}_{0+,0+}^q + \mathcal{F}_{0-,0-}^q = -C_F \sqrt{\frac{2}{N_c}} \frac{Q}{\xi} [T_s - T_u], \quad (13)$$

where N_c denotes the number of colors and $C_F = (N_c^2 - 1)/(2N_c)$ is the usual color factor. For convenience we only quote the sum over the quark helicities since this is what appears in (3). The denominators of the parton propagators read

$$T_s = \frac{1}{k_\perp^2 - \tau(\bar{x} - \xi)Q^2/(2\xi) - i\epsilon}, \quad T_u = \frac{1}{k_\perp^2 + \bar{\tau}(\bar{x} + \xi)Q^2/(2\xi) - i\epsilon}. \quad (14)$$

In both the quark and the gluon propagators we only retain k_\perp in the denominators of the parton propagators where it plays a crucial role. Its square competes with terms $\propto \tau(\bar{\tau})Q^2$ which become small in the end-point regions where either τ or $\bar{\tau}$ tends to zero.

While for longitudinally polarized vector mesons the quark transverse momentum is only needed for the suppression of the leading-twist contribution (the $\mathbf{k}_\perp \rightarrow 0$ limit), it plays an even more important role in the case of transverse polarization. In the collinear limit the spin wavefunction of the meson is $\Gamma_V \propto \not{\epsilon}' \not{\epsilon}_V(\pm 1)$ where ϵ_V denotes the polarization vector of the meson. It however leads to a vanishing contribution to the subprocess amplitude since the number of γ matrices in the Dirac trace³ that gives the hard scattering kernel, is odd (see Fig. 1)⁴. If one allows for quark transverse momenta a second term in the covariant spin wavefunction appears

$$\Delta\Gamma_V^\nu K_\nu = \frac{1}{\sqrt{2}M_V} \{ \not{\epsilon}' \not{\epsilon}_V, \gamma_\nu \} K^\nu, \quad (15)$$

for which the number of γ matrices in the Dirac trace is even. In eq. (15) M_V is a soft parameter of the order of the vector-meson mass, e.g. a typical constituent quark mass. The transverse momentum four-vector $K = [0, 0, \mathbf{k}_\perp]$ is suitably defined as the quark-antiquark relative momentum and represents one unit of orbital angular momentum in a covariant manner [1]. Expanding now the hard scattering kernel for transversely polarized vector mesons, one finds

$$\begin{aligned} \mathcal{F}_{+\lambda,+\lambda}^i(\xi, \bar{x}, \tau, Q^2, K) &= \mathcal{F}_{+\lambda,+\lambda}^i(\xi, \bar{x}, \tau, Q^2, \mathbf{k}_\perp^2) + \Delta\mathcal{F}_{+\lambda,+\lambda}^{i\nu}(\xi, \bar{x}, \tau, Q^2, \mathbf{k}_\perp^2) K_\nu \\ &+ \Delta\mathcal{F}_{+\lambda,+\lambda}^{i\nu\mu}(\xi, \bar{x}, \tau, Q^2, \mathbf{k}_\perp^2) K_\nu K_\mu + \dots \end{aligned} \quad (16)$$

Higher order terms in K_μ are neglected. In the spirit of the modified perturbative approach, the \mathbf{k}_\perp^2 terms in the propagator denominators are kept as has been done for the longitudinal amplitude. As already mentioned the first term in Eq. (16), generated by $\not{\epsilon}' \not{\epsilon}_V(\pm 1)$, vanishes as a consequence of the number of γ matrices in the trace and, evidently, the second one as well after integration on \mathbf{k}_\perp . Hence, the third term in (16) is the leading one for transversely polarized mesons and leads to the subprocess amplitudes

$$\mathcal{H}_{+\lambda,+\lambda}^{Vi} = -\frac{g_{\perp\mu\nu}}{2} \int d\tau \frac{d\mathbf{k}_\perp^2}{16\pi^2} \mathbf{k}_\perp^2 \Psi_{VT}(\tau, \mathbf{k}_\perp^2) \Delta\mathcal{F}_{+\lambda,+\lambda}^{i\nu\mu}, \quad (17)$$

in which g_\perp is the transverse metric tensor⁵. Note that the wavefunctions for longitudinally and transversely polarized vector mesons are different in general. The transverse amplitude is of order $\mathbf{k}_\perp^2/(M_V Q)$. Noting that $\langle \mathbf{k}_\perp^2 \rangle^{1/2}/M_V$ is of order unity, one realizes that the transverse amplitude is suppressed by

$$\mathcal{M}_{+\nu',+\nu} \propto \langle \mathbf{k}_\perp^2 \rangle^{1/2}/Q, \quad (18)$$

³ For the quark subprocess the trace includes the hadronic matrix element that defines the GPDs and is $\propto \gamma^+$ or $\propto \gamma^+ \gamma_5$.

⁴ Note that for longitudinally polarized vector mesons $\epsilon(0) \simeq q'/m_V$. Hence, the spin wavefunction is $\propto \not{q}'$ in this case up to mass corrections. A mass term $\propto m_V \not{\epsilon}_V$ in the spin wavefunction for transversely polarized vector mesons has been investigated [4]. In collinear approximation this term leads to an infrared singular twist-3 contribution. Such mass terms are neglected by us.

⁵ All its elements are zero except $g_\perp^{11} = g_\perp^{22} = -1$.

with respect to the one for longitudinally polarized vector mesons.

Working out the kernels, one finds after summation over the parton helicity that the kernel for transverse photon and meson polarization is obtained from the longitudinal one, Eq. (13), by the replacement

$$T_s - T_u \longrightarrow \frac{\mathbf{k}_\perp^2}{2} \frac{Q}{M_V} [T_s T_a - T_u T_b]. \quad (19)$$

The new propagator denominators read

$$T_a = \frac{1}{\tau Q^2 + \mathbf{k}_\perp^2}, \quad T_b = \frac{1}{\bar{\tau} Q^2 + \mathbf{k}_\perp^2}. \quad (20)$$

We note in passing that T_s and T_u represent the denominators of the gluon propagators in the LO subprocess $\gamma^* q \rightarrow V q$ while T_a and T_b belong to the quark propagators (see Fig. 1). With the help of partial fractioning ($i = s, u, j = a, b$)

$$T_i T_j = \frac{1}{\mathbf{k}_\perp^2} [c_i T_i + c_j T_j], \quad (21)$$

we can cast the Fourier transform of the transverse subprocess amplitude into exactly the same form as for the longitudinal one, Eq. (11). The kernel is then a linear combination of four Fourier transformed propagators.

The denominators of the parton propagators in (14) are either of the type

$$T_1 = \frac{1}{\mathbf{k}_\perp^2 + d_1 Q^2}, \quad (22)$$

or

$$T_2 = \frac{1}{\mathbf{k}_\perp^2 - d_2(\bar{x} \pm \xi)Q^2 - i\varepsilon}. \quad (23)$$

where $d_i \geq 0$. The Fourier transforms of these propagator terms can readily be obtained:

$$\begin{aligned} \hat{T}_1 &= \frac{1}{2\pi} K_0(\sqrt{d_1} bQ), \\ \hat{T}_2 &= \frac{1}{2\pi} K_0\left(\sqrt{d_2(\pm\xi - \bar{x})} bQ\right) \theta(\pm\xi - \bar{x}) \\ &\quad + \frac{i}{4} H_0^{(1)}\left(\sqrt{d_2(\bar{x} \pm \xi)} bQ\right) \theta(\bar{x} \pm \xi), \end{aligned} \quad (24)$$

where K_0 and $H_0^{(1)}$ are the zeroth order modified Bessel function of second kind and Hankel function, respectively.

3 The double distribution model

As in Refs. [1, 3] the GPDs are constructed from the PDFs with the help of double distributions [7, 8]. Since this construction is described in detail in our previous papers we will

only recapitulate a few basic elements of it. For details we refer to Refs. [1, 3]. The main advantage of this construction is the warranted polynomiality of the resulting GPDs and the correct forward limit $\xi, t \rightarrow 0$. It is well-known that, at low x , the parton distribution functions behave as powers δ_i of x . These powers are assumed to be generated by Regge poles [21, 22]. We generalize this behavior of the PDFs by assuming that the t dependence of the double distributions and hence the GPDs are also under control of Regge behavior. Linear Regge trajectories are assumed for small $-t$

$$\alpha_i = \alpha_i(0) + \alpha'_i t, \quad i = g, \text{sea}, \text{val}, \quad (25)$$

with $\delta_i = \alpha_i(0)$ for quarks and, as a consequence of the familiar definition of the gluon GPD which reduces to $\bar{x}g(\bar{x})$ in the forward limit, $\delta_g = \alpha_g(0) - 1$ for gluons. The trajectories are accompanied by Regge residues assumed to have an exponential t dependence with parameters b_i . The following ansatz for the double distributions associated with the GPDs H_i is therefore employed (cf. [3])

$$f_i(\beta, \alpha, t) = e^{b_i t} |\beta|^{-\alpha'_i t} h_i(\beta) \frac{\Gamma(2n_i + 2)}{2^{2n_i+1} \Gamma^2(n_i + 1)} \frac{[(1 - |\beta|)^2 - \alpha^2]^{n_i}}{(1 - |\beta|)^{2n_i+1}}, \quad (26)$$

where

$$\begin{aligned} h_g(\beta) &= |\beta|g(|\beta|) & n_g &= 2, \\ h_{\text{sea}}^q(\beta) &= q_{\text{sea}}(|\beta|) \text{sign}(\beta) & n_{\text{sea}} &= 2, \\ h_{\text{val}}^q(\beta) &= q_{\text{val}}(\beta) \Theta(\beta) & n_{\text{val}} &= 1. \end{aligned} \quad (27)$$

For the decomposition of the double distribution into valence and sea contribution we follow the procedure proposed in Ref. [12] and write

$$\begin{aligned} f_{\text{val}}^q(\beta, \alpha, t) &= [f^q(\beta, \alpha, t) + f^q(-\beta, \alpha, t)] \Theta(\beta), \\ f_{\text{sea}}^q(\beta, \alpha, t) &= f^q(\beta, \alpha, t) \Theta(\beta) - f^q(-\beta, \alpha, t) \Theta(-\beta). \end{aligned} \quad (28)$$

In the forward limit, $\xi, t \rightarrow 0$, this decomposition is conform to the usual definition of sea and valence quark PDFs.

The GPDs are related to the double distributions by the integral

$$H_i(\bar{x}, \xi, t) = \int_{-1}^1 d\beta \int_{-1+|\beta|}^{1-|\beta|} d\alpha \delta(\beta + \xi\alpha - \bar{x}) f_i(\beta, \alpha, t). \quad (29)$$

For convenience we employ an expansion of the PDFs ($\beta > 0$)

$$h_i(\beta) = \beta^{-\delta_i} (1 - \beta)^{2n_i+1} \sum_{j=0}^3 c_{ij} \beta^{j/2}, \quad (30)$$

which is particularly useful at low β and allows to perform the integral (29) term by term analytically; its use is also convenient if the integral (29) is carried out numerically. The

factor $(1 - \beta)^{2n_i+1}$ serves for canceling the corresponding factor in Eq. (26) and has the additional welcome feature of roughly accounting for the $\beta \rightarrow 1$ behavior of the PDFs. The ansatz (30) results in a corresponding expansion of the GPDs

$$H_i(\bar{x}, \xi, t) = e^{b_i t} \sum_{j=0}^3 c_{ij} H_{ij}(\bar{x}, \xi, t). \quad (31)$$

The definition of the GPDs is completed by the relations

$$H^g(-\bar{x}, \xi, t) = H^g(\bar{x}, \xi, t), \quad H_{\text{sea}}^q(-\bar{x}, \xi, t) = -H_{\text{sea}}^q(\bar{x}, \xi, t), \quad (32)$$

and

$$H_{\text{val}}^q(\bar{x}, \xi, t) = 0 \quad -1 \leq \bar{x} < -\xi. \quad (33)$$

In Eq. (29) so-called D terms for the gluons and the flavor singlet quark combination are ignored [23]. The D terms ensure the appearance of the highest powers of the skewness in the moments of the GPDs. They only contribute to the less important real parts of the amplitudes since their support is the region $-\xi < x < \xi$. The corresponding imaginary parts are related to the GPDs at $\bar{x} = \xi(1 + 2\mathbf{k}_\perp^2/(\tau Q^2))$ which lies outside the support of the D terms. We take this in vindication of neglecting the D terms.

For the expansions of the PDFs we will use the same parameters as in Ref. [3] with the exception of a little change. ZEUS [24] now provides data on the cross section for ρ production up to $Q^2 = 100 \text{ GeV}^2$ whereas in [3] the fits to the CTEQ6M PDFs were made for $Q^2 \leq 40 \text{ GeV}^2$. As one may check, the CTEQ6M gluon and sea quark PDFs are not well described above 40 GeV^2 by the expansion quoted in Ref. [3] but the addition of a L^2 -term to δ_g

$$\delta_g = 0.10 + 0.06L - 0.0027 L^2, \quad (34)$$

improves the fits to the gluon and sea quark PDFs considerably as can be seen from Fig. 2 ($L = \ln Q^2/Q_0^2$, $Q_0^2 = 4 \text{ GeV}^2$). The L^2 term is irrelevant below 40 GeV^2 . The valence quarks are not needed at high Q^2 . The parameters of the expansions (30) are quoted in Tab. 1. We stress that with the exception of δ_g , they are identical to those used in Ref. [3]. In the quoted ranges of Q^2 and β the fits to the PDFs agree very well with the CTEQ6M solution; they are always well inside the band of Hessian errors quoted in [6]. Larger values of β are irrelevant to us since the region $0.5 \lesssim \beta$ affects the real parts of the amplitudes only marginally; the contributions are less than 0.5%. From the fitted PDFs the GPDs are evaluated with the help of Eq. (29).

In an attempt to keep the GPD model simple we assume

$$H_{\text{sea}}^u = H_{\text{sea}}^d = \kappa_s H_{\text{sea}}^s, \quad (35)$$

where the flavor symmetry breaking factor is parameterized as

$$\kappa_s = 1 + 0.68/(1 + 0.52 \ln Q^2/Q_0^2), \quad (36)$$

as obtained from a fit to the CTEQ6M PDFs.

	gluon	strange	u_{val}	d_{val}
δ	Eq. (34)	$1 + \delta_g$	0.48	0.48
α'	0.15 GeV^{-2}	0.15 GeV^{-2}	0.9 GeV^{-2}	0.9 GeV^{-2}
c_0	$2.23 + 0.362 L$	$0.123 + 0.0003 L$	$1.52 + 0.248 L$	$0.76 + 0.248 L$
c_1	$5.43 - 7.00 L$	$-0.327 - 0.004 L$	$2.88 - 0.940 L$	$3.11 - 1.36 L$
c_2	$-34.0 + 22.5 L$	$0.692 - 0.068 L$	$-0.095 L$	$-3.99 + 1.15 L$
c_3	$40.6 - 21.6 L$	$-0.486 + 0.038 L$	0	0

Table 1: The parameters appearing in the expansion (30) of the PDFs ($L = \ln Q^2/Q_0^2$, $Q_0^2 = 4 \text{ GeV}^2$).

As in Ref. [3] we take for the slope of the gluon trajectory the value $\alpha'_g = 0.15 \text{ GeV}^{-2}$. Since the sea quark PDFs are mainly driven by evolution for $Q^2 \gtrsim 4 \text{ GeV}^2$ it is assumed that $\alpha_{\text{sea}}(t) = \alpha_g(t)$. A standard trajectory is adopted for the valence quark Regge pole - $\alpha_{\text{val}}(t) = 0.48 + 0.9 \text{ GeV}^{-2} t$. The parameter of the gluon residue is fixed from a fit against the HERA data for ρ [25] and ϕ production [26]:

$$b_g = b_{\text{sea}} = 2.58 \text{ GeV}^{-2} + 0.25 \text{ GeV}^{-2} \ln \frac{m^2}{Q^2 + m^2}, \quad (37)$$

The ρ and ϕ slopes of the cross sections practically fall together at HERA energies; there are only minor differences at low Q^2 . The parameter b_g given in (37) leads to a t dependence of the differential cross section in perfect agreement with the recent ZEUS data [24]. The parameter of the valence quark residue is taken to be zero. This is in accord with the findings of the nucleon form factor analysis proposed in Ref. [9] in which the zero-skewness GPDs have been determined.

We emphasize that the evolution of the GPDs is taken into account by us only approximately through the evolution of the PDFs. This is reasonable since at low ξ the imaginary part of the gluon (and sea quark) contribution dominates which is $\propto H^g(\xi, \xi, t)$ and therefore approximately equals $2\xi g(2\xi)$ at low ξ (see, for instance [1]). Its real part as well as the valence quark contributions are only of importance near 4 GeV^2 , the initial value of the evolution. The very time-consuming numerical integration on x , \mathbf{b} and τ forces us to use this approximative treatment of the evolution. In order to demonstrate the quality of our approximation we compare in Fig. 2 the gluon GPD at 40 GeV^2 either obtained from our approximation or from evolving H^g from the initial scale of 4 GeV^2 using the evolution code developed by Vinnikov [27]. Only minor differences are to be noticed.




Figure 2: Left: The CTEQ6 gluon PDF at $Q^2 = 100 \text{ GeV}^2$ as compared to our fit (solid line). The band indicates the Hessian errors of the PDFs. Right: The gluon GPD H^g at $\xi = 0.01$ and $t = 0$. Solid line: H^g at the initial scale 4 GeV^2 . Dash-dotted (dashed) line: at the scale of 40 GeV^2 evolved with the Vinnikov code [27] (approximate evolution).

4 Results on cross sections

Before we present our results obtained from the handbag approach we have to specify the meson wave functions used in the evaluation of the amplitudes. As in Refs. [1, 3] we will take Gaussian wave functions ($j = L, T$)

$$\Psi_{Vj}(\tau, \mathbf{k}_\perp) = 8\pi^2 \sqrt{2N_c f_{Vj}(\mu_F) a_{Vj}^2} \left[1 + B_2^{Vj}(\mu_F) C_2^{3/2} (2\tau - 1) \right] \exp[-a_{Vj}^2 \mathbf{k}_\perp^2 / (\tau \bar{\tau})]. \quad (38)$$

Transverse momentum integration leads to the meson distribution amplitudes for which we allow for the second Gegenbauer moment besides the asymptotic form. The meson decay constants for longitudinally polarized vector mesons are known from the electronic decays of the vector mesons while those for transversely polarized mesons are taken from QCD sum rules [28]. In contrast to the decay constants f_{VL} the latter ones are scale dependent

$$f_{VT}(\mu_F) = f_{VT}(\mu_0) \left(\frac{\alpha_s(\mu_F)}{\alpha_s(\mu_0)} \right)^{4/27}. \quad (39)$$

Note that the decay constants for transversely polarized vector mesons always appear in the combination f_{VT}/M_V , i.e. there is only one independent parameter in practice. In fact we use a typical constituent mass of 300 MeV for M_V and the QCD sum rule result [28] $f_{VT}/f_{VL} \simeq 0.8$ at the scale of $\mu_0 = 1 \text{ GeV}$. The Gegenbauer coefficients B_2^{VL} have been found to be zero in the analysis of the longitudinal cross section [3]. Those for the transverse case are fitted to the data on the σ_T or the cross section ratio $R = \sigma_L/\sigma_T$. The

parameter	ρ_L	ρ_T	ϕ_L	ϕ_T
$f_V[\text{MeV}]$	209	167	221	177
$a_V[\text{GeV}^{-1}]$	0.75	1.0	0.70	0.95
B_2^V	0.0	0.10	0.0	0.10

Table 2: The parameters appearing in the wavefunction (38), quoted at the scale $\mu_0 = 1 \text{ GeV}$.

Gegenbauer coefficients are scale dependent

$$B_2^{Vj}(\mu_F) = B_2^{Vj}(\mu_0) \left(\frac{\alpha_s(\mu_F)}{\alpha_s(\mu_0)} \right)^{\gamma_{2j}}, \quad (40)$$

where $\gamma_{2L} = 50/81$ and $\gamma_{2T} = 40/81$ [29]. Finally, the transverse size parameters a_{Vj} in (38) are either fitted to σ_L or to σ_T depending on the polarization of the vector meson. The values for the various parameters are compiled in Tab. 2. Those for the longitudinal case are identical to the parameters used in [3].

The assessment of the theoretical uncertainties deserves special considerations. The results on cross sections (and other observables) are subject to parametric errors. The main uncertainties stem from the Hessian errors of the set of CTEQ6 PDFs. Since for the longitudinal cross section the parameters of the corresponding wave functions are adjusted such that good agreement between the data on σ_L and the handbag results is achieved, there is no substantial additional uncertainties from the longitudinal wavefunction. Results for σ_L evaluated from sets of PDFs other than CTEQ6 also fall into the error bands in most cases (an exception is set for instance by the PDFs determined in Ref. [30, 31]) provided these PDFs are treated in analogy to the CTEQ6M set, i.e. they are fitted to the expansion (30) by forcing them to behave Regge-like with powers δ_i as described above, and, if necessary, readjusting the transverse size parameters. The uncertainties in the ratio, R , of the longitudinal and transverse cross sections mainly arise from the uncertainties of the wavefunctions for transversely polarized vector mesons (i.e. from the Gegenbauer coefficients, the transverse size parameters and from the ratio f_{VT}/M_V). The errors due to those of the GPDs or PDFs which are mainly responsible for the theoretical uncertainties of the cross sections, cancel to a large extent in the ratio.

The results for the longitudinal cross sections are the same as in Ref. [3]. We refrain from showing them again. The experimental data on the cross section ratio R are customarily determined from the measured SDME r_{00}^{04} by the relation

$$R = \frac{\sigma_L}{\sigma_T} = \frac{1}{\varepsilon} \frac{r_{00}^{04}}{1 - r_{00}^{04}}. \quad (41)$$

The SDME in (41) is understood to be integrated over the full range of t available in a

given experiment ⁶. The theoretical and experimental results on the ratio R are compared in Figs. 3 and 4. In general we achieve very good agreement with experiment in particular with regard to the theoretical uncertainties displayed as shaded bands in the plots.

The ratio R is mildly energy dependent for W larger than about 10 GeV while, for lower energies, it exhibits a somewhat stronger energy dependence, in particular at larger values of Q^2 , see Fig. 4. This implies differences in the energy dependences of σ_L and σ_T which can be traced back to the different hard scattering kernels (see e.g. Eq. (19)) and the varying wavefunctions for longitudinally and transversely polarized vector mesons, in particular to the different values of the r.m.s. \mathbf{k}_\perp ($\sim 1/a_V$). An energy dependence of R seems to be indicated by the preliminary HERMES [35] and COMPASS [38] data although confirmation of this observation is demanded. Most of the older experiments have rather large errors such that a definite conclusion on a possible W dependence cannot be drawn at present. Unfortunately, for ϕ production there is only one low energy data point available [37] and this point is measured at the very low value of $Q^2 = 2.6 \text{ GeV}^2$ which lies outside the range where the handbag approach, in its present form, can be trusted. We note that for ϕ production the energy dependence of R is even milder than for the case of the ρ ; the results for $W = 10 \text{ GeV}$ practically fall together with those at 75 GeV.

In Fig. 5 the handbag results on the ρ and ϕ cross sections are compared to experiment. Again good agreement with the H1 [25, 32] and ZEUS [24, 26, 34] data is to be observed in a large range of Q^2 . The leading-twist contribution to these cross sections, i.e. σ_L evaluated in collinear approximation, is also shown. Although the leading-twist contribution approaches the experimental cross section with increasing Q^2 there is still a small difference of about 1.5σ between both at $Q^2 = 100 \text{ GeV}^2$ for ρ production. Note that even at that value of Q^2 the transverse cross section which is included in the full one and also represents a power correction to the leading-twist result, is not negligible; it amounts to about 10%. The leading-twist contribution is about 20% larger than the one obtained within the modified perturbative approach at $Q^2 = 100 \text{ GeV}^2$, i.e. the corrections due the quark transverse momentum have not yet disappeared completely. In Fig. 6 the energy dependence of the ρ cross section at a set of Q^2 values is displayed. Within errors agreement is to be seen with the ZEUS data [24]. This is not a surprise since the power δ_g , related to the intercept of the gluonic Regge intercept, is fixed by the energy dependence of the HERA data on the cross sections, see Ref. [3]. As σ_L , see [3], the full cross section at HERA energies is dominated by the gluon contribution although the sea quarks are not negligible. Including the interference with the gluon the sea quarks contribute about 25% in the case of the ϕ and 40% in the case of the ρ at $Q^2 = 4 \text{ GeV}^2$. The larger sea-quark contribution in the latter case is due to the flavor-symmetry breaking factor κ_s (36). Flavor symmetry breaking in the sea is important for the ratio of the ϕ and ρ cross sections, see Ref. [3]. Neglecting the sea quarks or assuming a flavor symmetric sea leads to an incorrect $\phi - \rho$ ratio. Going to energies lower than about 10 GeV the valence quark contributions gradually become perceptible for ρ production. At, say, 5 GeV the valence quarks are responsible for about 40% of the

⁶ The cross sections σ_L and σ_T have been obtained from integrating the differential cross sections over that range of t .




Figure 3: The ratio of longitudinal and transverse cross sections for ρ (left) and ϕ (right) production versus Q^2 at $W = 90$ GeV and 75 GeV, respectively. Data taken from H1 [25, 32, 33] (solid squares) and ZEUS [26, 34] (open squares), respectively. The recent ZEUS data [24] are shown as open triangles. The solid lines represent the handbag results with the shaded bands indicating the uncertainties of the predictions.




Figure 4: Handbag results for R on ρ (left) and ϕ (right) production at $W = 75(10, 5)$ GeV shown as solid (dash-dotted, dashed) line. Data taken from Refs. [25, 26, 34]. Preliminary data from HERMES [35, 37] (solid circle) and COMPASS [38] (diamond). The error bands are only shown at $W = 5$ GeV. For further notation refer to Fig. 3.

cross section. We stress that the three contributions have to be added coherently. There are substantial interference terms which increase the cross sections markedly. For instance, at $W = 5 \text{ GeV}$ and $Q^2 = 4 \text{ GeV}^2$ the ρ cross section is doubled by the interference terms.

A remark concerning the t dependence of the differential cross sections is in order. As we pointed out in [3] the cross sections drop exponentially with t to a very good approximation. Their slopes are approximately given by $2b_g$ (see (37)) plus a contribution from the gluonic Regge trajectory (see Eq. (26) and Ref. [3]). At large W the slopes of the differential cross sections for longitudinally and transversely polarized photons are the same while, at low W , the valence quarks generate small differences. Consequently, the ratio R is nearly t independent a fact that is in agreement with the ZEUS measurement [24].

Finally, we check the theoretical consistency of the modified perturbative approach. Consistency is meant in the sense that the bulk of the perturbative contributions should be accumulated in regions where the strong coupling α_s is sufficiently small. To find out whether or not this is the case, we set the integrand in (11) equal to zero in those regions where $\alpha_s(\mu_R) > \alpha_s^{\text{crit}}$, and evaluate, say, the longitudinal cross section for ρ production as a function of α_s^{crit} . Consulting Fig. 6 where the accumulation profile is shown, one observes that almost the entire result is accumulated in a comparatively narrow region of α_s around 0.4 at $Q^2 = 4 \text{ GeV}^2$ and 0.25 at 40 GeV^2 . Thus the effective renormalization scales in these two cases are about 1.5 and 15 GeV^2 , respectively. Hence, our results on cross sections are theoretically self-consistent with regard to the above-mentioned criterion. Contributions from the end-point regions where the momentum fraction τ tends either to zero or to one and where, in collinear approximation, the renormalization scale becomes very small, are sufficiently suppressed.

5 Results on spin density matrix elements

In a number of experiments [24, 25, 32, 38, 39] the SDMEs have been extracted from the measured decay angular distributions of the vector mesons. The formalism of the SDMEs, i.e. their relations to the amplitudes for the process $\gamma^*p \rightarrow Vp$, has been developed by Schilling and Wolf [40] long-time ago. This work has recently been repeated and extended to the case of a transversely polarized target proton by Diehl [41]. Since in the experimental papers the notation of Ref. [40] is used throughout, we will adhere to it here in order to facilitate comparison.

The SDMEs are given by properly normalized bilinear combinations of $\gamma^*p \rightarrow Vp$ amplitudes. Due to the symmetry relations (9) and (10) there are no interference terms between \mathcal{M}^N and \mathcal{M}^U in the case of unpolarized electroproduction of vector mesons. The normalizations read

$$\begin{aligned} N_L &= 2 \sum_{\nu'} |\mathcal{M}_{0\nu',0+}^N|^2, \\ N_T &= 2 \sum_{\nu'} \left[|\mathcal{M}_{+\nu',++}^N|^2 + |\mathcal{M}_{+\nu',++}^U|^2 \right], \end{aligned} \quad (42)$$




Figure 5: The Q^2 -dependence of the cross sections for ρ (left) and ϕ (right) production at $W = 90$ (ρ) and 75 GeV (ρ, ϕ). Data are taken from ZEUS [24, 26, 34] and H1 [25, 32]. For ρ production at $W = 75$ GeV the data and theoretical results are divided by 10 for the ease of legibility. The dashed lines represent the leading-twist contribution. For further notations refer to Fig. 3.




Figure 6: Left: The cross section for ρ production versus W at a set of Q^2 values. The data are taken from ZEUS [24]. For further notations refer to Fig. 3. The error band at 6 GeV^2 is omitted for the ease of legibility. Right: The accumulation profile of the amplitude in dependence of α_c^{crit} at $Q^2 = 4$ (solid line) and 40 GeV^2 (dashed line) at $W = 75 \text{ GeV}$.

where helicity-flip $\gamma^* \rightarrow V$ transitions are neglected as we do throughout this work. Up to a phase space factor, see Eq. (7), the normalizations are the differential cross sections for longitudinally and transversely polarized vector mesons. Since the SDMEs also provide a possibility to study the role of \widetilde{H} we do not ignore the contributions from the corresponding amplitude $\mathcal{M}_{+\nu',++}^U$ in (42) for later reference.

Obviously, the SDMEs are functions of Q^2, t and W . Due to limitations in statistics it is not possible to measure the SDMEs as a functions of the three variables. Thus, frequently the SDMEs are presented as functions of one variable for average values of the other variables. For instance the SDMEs are quoted as a function of Q^2 for an average value of W and integrated over all t available in a given experiment. Only in this case, and this is an important one to which we will mainly refer to in the following, the normalizations N_L and N_T refer to the respective integrated cross sections σ_L and σ_T up to the phase-space factor and eventually neglected suppressed amplitudes. If for instance the SDMEs are presented as a function of t the normalization represents differential cross sections at a fixed value of t but integrated over certain ranges of Q^2 and W .

The SDMEs allow for a separation of the absolute values of the two amplitudes, $\mathcal{M}_{0+,0+}^N$ and $\mathcal{M}_{++,++}^N$, and for a test of their relative phases. Whether or not it is justified to neglect the other amplitudes can be examined as well. The SDMEs r_{00}^{04} , r_{1-1}^1 and $\text{Im } r_{1-1}^2$ are related to the cross section ratio R . In terms of the amplitudes we are investigating these SDMEs read

$$\begin{aligned} r_{00}^{04} &= \frac{2\varepsilon}{N_T + \varepsilon N_L} \sum_{\nu'} |\mathcal{M}_{0\nu',0+}^N|^2, \\ r_{1-1}^1 &= -\text{Im} r_{1-1}^2 = \frac{1}{N_T + \varepsilon N_L} \sum_{\nu'} [|\mathcal{M}_{+\nu',++}^N|^2 - |\mathcal{M}_{+\nu',++}^U|^2]. \end{aligned} \quad (43)$$

Neglecting the amplitude \mathcal{M}^U too these expressions simplify to

$$1 - r_{00}^{04} = 2r_{1-1}^1 = -2\text{Im } r_{1-1}^2 = \frac{1}{1 + \varepsilon R}. \quad (44)$$

In Figs. 7, 8 and 9 this prediction of our handbag approach is compared to experiment. The agreement is in general good within errors although with occasional exceptions. Thus, for ρ production, r_{1-1}^1 and $\text{Im } r_{1-1}^2$ are not perfectly reproduced. On the other hand, their sum, measuring the double flip transitions $\gamma_T^* \rightarrow V_{-T}$, is nicely in agreement with zero. A further check is provided by the t dependence of the SDMEs. Since in the handbag approach the t dependence solely comes from the GPDs and these are identical for the two amplitudes the above SDMEs should be nearly flat in t which is indeed the case experimentally within errors [24, 25]. Hence, the above three SDMEs are consistent with our assumptions and do not provide a significant signal for helicity-flip amplitudes or contributions from \widetilde{H} . The W dependence of the handbag predictions is displayed in Fig. 8. It is evidently very mild. In particular the results for the SDMEs at $W = 75$ and 90 GeV fall practically together.

After having checked that the absolute values of our two amplitudes are in fair agreement with experiment, we now turn to their relative phase δ_{LT} . In terms of our two

amplitudes the SDMEs relevant for the verification of the phase, read

$$\begin{aligned} \text{Re } r_{10}^5 &= -\text{Im } r_{10}^6 = \frac{1}{\sqrt{2}} \frac{1}{N_T + \varepsilon N_L} \text{Re} [\mathcal{M}_{++,++}^N \mathcal{M}_{0+,0+}^{N*}] \\ &= \frac{1}{2\sqrt{2}} \frac{\sqrt{R}}{1 + \varepsilon R} \cos \delta_{LT}. \end{aligned} \quad (45)$$

Predictions for the SDMEs in the case of ϕ production are shown in Figs. 9 and 10. Fair agreement with experiments can be seen. For ρ production, on the other hand, a conflict is to be noted, see Figs. 7 and 8. The data [24, 25, 35] require a rather large phase although with strong fluctuations ($10 - 30^\circ$) while the handbag approach provides only a small value for it (e.g. $\delta_{LT} = 3.1^\circ$ at $W = 5$ GeV and $Q^2 = 3$ GeV²). Whether our model for the $\gamma_T^* \rightarrow V_T$ amplitude which represents a power correction to the leading $\gamma_L^* \rightarrow V_L$ amplitude, is inadequate for this detail needs further investigation. However, that the sum $\text{Re } r_{10}^5 + \text{Im } r_{10}^6$ amounts to only 1% of the corresponding difference of these SDME makes it clear that the neglected helicity flip $\gamma^* \rightarrow V$ transitions in (45) are not responsible for the observed conflict.

The HERMES collaboration has also measured the decay angular distribution of the ρ in the case of a longitudinally polarized lepton beam. These measurements which are not yet analyzed, will provide data on other SDMEs [40]. For instance, the SDMEs $\text{Im } r_{10}^7$ and $\text{Re } r_{10}^8$ also measure the phase δ_{LT} introduced in (45)

$$\text{Im } r_{10}^7 = \text{Re } r_{10}^8 = \frac{1}{2\sqrt{2}} \frac{\sqrt{R}}{1 + \varepsilon R} \sin \delta_{LT}. \quad (46)$$

Predictions for these SDMEs are shown in Figs. 11. The other polarized SDMEs are only sensitive to the suppressed amplitudes, see Tab. 3.

In principle also the helicity flip $\gamma^* \rightarrow V$ transitions can be calculated in the proposed handbag approach as well. But these transitions are strongly suppressed with respect to the amplitude $\mathcal{M}_{0+,0+}$ as the amplitude for $\gamma_T^* \rightarrow V_T$ transitions, see (18). In fact,

$$\begin{aligned} \gamma_T^* \rightarrow V_L &\propto \frac{\sqrt{-t}}{Q} \\ \gamma_L^* \rightarrow V_T &\propto \frac{\sqrt{-t} \langle \mathbf{k}_\perp^2 \rangle^{1/2}}{Q^2} \\ \gamma_T^* \rightarrow V_{-T} &\propto \frac{-t \langle \mathbf{k}_\perp^2 \rangle^{1/2}}{Q^3}, \end{aligned} \quad (47)$$

where the powers of $\sqrt{-t}/Q$ follow from angular momentum conservation and the factors $\langle \mathbf{k}_\perp^2 \rangle^{1/2}/Q$ from the treatment of transversely polarized vector meson, see (18). The suppression factors given in (47) engender the behavior of the SDMEs for helicity flip $\gamma^* \rightarrow V$ transitions listed in Tab. 3. The most important helicity-flip amplitude is the one for $\gamma_T^* \rightarrow V_L$ transitions. It is most clearly seen in r_{00}^5 as an interference term with the dominant longitudinal amplitude. It is definitely non-zero, of the order of 0.1 for ρ and ϕ




Figure 7: The SDMEs for the ρ meson versus Q^2 (in GeV^2) at $W = 75$ GeV. Data taken from [24, 25, 39]. Note the different scales at the axis of ordinates. For further notations refer to Fig. 3.




Figure 8: SDMEs for the ρ meson versus Q^2 (in GeV^2) at $W = 5, 10$ and 75 GeV (dashed, dash-dotted and solid line, respectively). Preliminary data taken from HERMES [35] (solid circles) and COMPASS [38] (diamonds). Error bands only shown for $W = 5$ GeV. For further notations refer to Figs. 3 and 4.




Figure 9: SDMEs for the ϕ meson versus Q^2 (in GeV^2) at $W = 75$ GeV. Data taken from [26, 32]. For further notations refer to Fig. 3.




Figure 10: SDMEs for the ϕ meson versus Q^2 (in GeV^2) at $W = 5$ GeV.




Figure 11: The SDMEs $\text{Im } r_{10}^7$ and $\text{Re } r_{10}^8$ for ρ production versus Q^2 at $W = 5$ GeV. For further notations refer to Fig. 3.

SDME	amplitudes	order
r_{00}^5, r_{00}^8	$(\gamma_T^* \rightarrow V_L)(\gamma_L^* \rightarrow V_L)$	$\sqrt{-t}/Q$
$\text{Re } r_{10}^{04}, \text{Re } r_{10}^1, \text{Im } r_{10}^2, \text{Im } r_{10}^3$	$(\gamma_T^* \rightarrow V_L)(\gamma_T^* \rightarrow V_T)$	$\sqrt{-t}\langle \mathbf{k}_\perp^2 \rangle^{1/2}/Q^2$
r_{00}^1	$ (\gamma_T^* \rightarrow V_L) ^2$	$-t/Q^2$
$r_{11}^5, r_{1-1}^5, \text{Im } r_{1-1}^6$	$(\gamma_L^* \rightarrow V_T)(\gamma_T^* \rightarrow V_T)$	$\sqrt{-t}\langle \mathbf{k}_\perp^2 \rangle/Q^3$
$r_{11}^8, r_{1-1}^8, \text{Im } r_{1-1}^7$	$(\gamma_L^* \rightarrow V_T)(\gamma_T^* \rightarrow V_T)$	$\sqrt{-t}\langle \mathbf{k}_\perp^2 \rangle/Q^3$
$r_{1-1}^{04}, r_{11}^1, \text{Im } r_{1-1}^3$	$(\gamma_T^* \rightarrow V_{-T})(\gamma_T^* \rightarrow V_T)$	$-t\langle \mathbf{k}_\perp^2 \rangle/Q^4$

Table 3: SDMEs controlled by helicity flip $\gamma^* \rightarrow V$ transitions.

production in experiment [24, 25, 35]. At least HERMES [35, 36] observes a t dependence for it in agreement with expectation ($\propto \sqrt{-t}$). The contribution of the $\gamma_T^* \rightarrow V_L$ amplitude to the cross sections, R and r_{00}^{04} is at the percent level and can safely be neglected. The other SDMEs related to $\gamma_T^* \rightarrow V_L$ transitions are experimentally compatible with zero within errors. The only exception is to be seen in the recent high statistics ZEUS data [24] where $\text{Im } r_{10}^2$ and $\text{Re } r_{10}^1$ have very small but non-zero for values of Q^2 less than 10 GeV². It would be interesting to see whether their t dependencies are in agreement with expectation. For the remaining SDMEs being related to $\gamma_L^* \rightarrow V_T$ and $\gamma_T^* \rightarrow V_{-T}$ transitions there is no significant deviation from zero at large Q^2 experimentally. In view of these remarks it is fair to conclude that the neglect of the helicity flip $\gamma^* \rightarrow V$ transitions in the cross sections for $Q^2 \geq 3$ GeV² is justified.

6 The role of \widetilde{H}

The expression for the amplitude \mathcal{M}^U is the same as that for \mathcal{M}^N given in (3) except that the sum of the subprocess amplitudes is to be replaced by their difference

$$\mathcal{M}_{\mu+, \mu+}^{Ui}(V) = \frac{e}{2} \sum_a e_a \mathcal{C}_V^a \int_{\bar{x}_i}^1 d\bar{x} \left[\mathcal{H}_{\mu+, \mu+}^{Vi} - \mathcal{H}_{\mu-, \mu-}^{Vi} \right] \widetilde{H}^i(\bar{x}, \xi, t). \quad (48)$$

The superposition of the various quark and gluon contributions is identical to that for the amplitude \mathcal{M}^N . The unnatural parity amplitudes satisfy the symmetry relation (10). It is evident from (48) that parity conservation leads to a vanishing longitudinal amplitude $\mathcal{M}_{0+, 0+}^U$. The transverse subprocess amplitude is the same as in (17) and (19) except that in the latter equation a plus sign occurs between $T_s T_a$ and $T_u T_b$ now.

The GPDs \widetilde{H} are again modeled by the double distribution ansatz (26). With regard to the symmetry properties of \widetilde{H} the functions \tilde{h}_i now take the form

$$\begin{aligned} \tilde{h}_g(\beta) &= |\beta| \Delta g(|\beta|) \text{sign}(\beta), \\ \tilde{h}_{\text{sea}}^q(\beta) &= \Delta q_{\text{sea}}(|\beta|), \\ \tilde{h}_{\text{val}}^q(\beta) &= \Delta q_{\text{val}}(\beta) \Theta(\beta). \end{aligned} \quad (49)$$

For the powers n_i in the double distribution ansatz (26) the same values are taken as for the GPDs H^i , see (27). The decomposition of the double distribution into valence and sea contributions is made by [12]

$$\begin{aligned} \tilde{f}_{\text{val}}^q(\beta, \alpha, t) &= [\tilde{f}^q(\beta, \alpha, t) - \tilde{f}^q(-\beta, \alpha, t)] \Theta(\beta), \\ \tilde{f}_{\text{sea}}^q(\beta, \alpha, t) &= \tilde{f}^q(\beta, \alpha, t) \Theta(\beta) + \tilde{f}^q(-\beta, \alpha, t) \Theta(-\beta). \end{aligned} \quad (50)$$

The double distribution ansatz for \widetilde{H}^g is incomplete because in moments of this GPD the highest power of ξ are lacking which leads to difficulties with the analytic properties of the amplitudes [42]. We ignore this problem here since the contributions from \widetilde{H}^g seem to be unimportant.

The required polarized parton distributions are taken from Ref. [43] and expanded according to

$$\tilde{h}_i(\beta) = \beta^{-\tilde{\delta}_i} (1 - \beta)^{2n_i+1} \sum_{j=0}^3 \tilde{c}_{ij} \beta^j, \quad (51)$$

using only integer powers. The resulting expansion parameters \tilde{c}_i and $\tilde{\delta}_i$ are quoted in Tab. 4. It is expected that the a_1 Regge trajectory controls the low- x behavior of the polarized valence quark PDFs. Since there are no recurrences of the $a_1(1260)$ we are forced to assume the standard value of 0.9 GeV^{-2} for the slope of the trajectory⁷. Combining this with the spin of the a_1 , we obtain $\alpha_{a_1}(0) \simeq -0.36$ for the intercept. Such a low value is however

⁷ Accepting exchange degeneracy for the a_1 and $\eta_2(1617)$ trajectories the slope of the trajectory is fixed by the meson spectrum and is indeed 0.9 GeV^{-2} .

	Δu_{val}	Δd_{val}	e_{val}^u	e_{val}^d
$\tilde{\delta}$	0.48	0.48	0.48	0.48
\tilde{c}_0	$0.61 + 0.033 L$	$-0.320 - 0.040 L$	2.204	-3.114
\tilde{c}_1	$0.410 - 0.377 L$	$-1.427 - 0.176 L$	-2.204	8.096
\tilde{c}_2	$5.10 - 1.21 L$	$0.692 - 0.068 L$	0.0	-6.477
\tilde{c}_3	0.0	0.0	0.0	1.295

Table 4: The parameters appearing in the expansion (30) of the polarized PDFs and the forward limits of E^a . The latter are taken from Ref. [9]. The expansion (30) provides a fit to the Blümlein-Böttcher PDFs [43] in the range $10^{-2} \leq \beta \leq 0.5$ and $4 \text{ GeV}^2 \leq Q^2 \leq 40 \text{ GeV}^2$. The powers $\tilde{\delta}$ are kept fixed in the fits to the PDFs.

in conflict with the small- x behavior⁸ of the polarized valence quark PDFs determined in Ref. [43], for which the power is rather about 0.7. As a compromise we therefore take the standard value of 0.48 for it. For the slope of the Regge trajectory we again take the value of 0.9 GeV^2 and for its residue $\tilde{b}_{\text{val}} = 0$.

The GPDs \tilde{H}^i are obtained from the functions f_i by the integral (29). They satisfy the relations

$$\tilde{H}^g(-\bar{x}, \xi, t) = -\tilde{H}^g(\bar{x}, \xi, t), \quad \tilde{H}_{\text{sea}}^q(-\bar{x}, \xi, t) = \tilde{H}_{\text{sea}}^q(\bar{x}, \xi, t), \quad (52)$$

and

$$\tilde{H}_{\text{val}}^q(\bar{x}, \xi, t) = 0 \quad -1 \leq \bar{x} < -\xi. \quad (53)$$

We checked that our proposed GPDs \tilde{H}_{val} are in agreement with the data on the axial form factor for $-t \lesssim 0.6 \text{ GeV}^2$ and with the low $-t$ (low x) behavior of \tilde{H} determined in [9]. In constrast to the situation for H , \tilde{H}_{val}^u and \tilde{H}_{val}^d have opposite signs as a consequence of the behavior of the polarized PDFs. The lowest moments of the latter are known from β decays (see, for instance, Ref. [43]). The usual assumption of a smooth behavior of the PDFs without a change of sign, leads to opposite signs of Δu_{val} and Δd_{val} . As our numerical studies reveal the gluon and sea quark contributions to \mathcal{M}^U are very small and compensate each other to a large extent since the gluonic and sea quark GPDs \tilde{H} have opposite signs. Their combined contributions are practically negligible. This is the reason why we quote only the expansion parameters of \tilde{h}_i for the valence quarks in Tab. 4.

Neglecting as in the preceding sections proton helicity flips and photon-meson transitions other than $L \rightarrow L$ and $T \rightarrow T$, one can project out the unnatural parity amplitude for $T \rightarrow T$ transitions from a particular combination of SDMEs (see Eq. (43))

$$U = \frac{1}{2} \left[1 - r_{00}^{04} - 2r_{1-1}^1 \right]$$

⁸ Whether this is a consequence of lack of low- x data in the PDF analysis or due to disregarded high-lying Regge cuts, is unknown at present.




Figure 12: Left: The ratio of σ_U and σ for ρ production versus Q^2 at $W = 5$ GeV. Data taken from HERMES [35]. The solid (dashed, dash-dotted) line represents our estimate (obtained with $e_u H_{\text{val}}^u + e_d H_{\text{val}}^d$, with $e_u H_{\text{val}}^u - e_d H_{\text{val}}^d$), see text. Right: The helicity correlation A_{LL} for ρ production at $W = 5$ (10) GeV dashed (dash-dotted) line. Data taken from COMPASS [48] and HERMES [49]. For other notations cf. to Figs. 3 and 4.

$$= \frac{2}{N_T + \varepsilon N_L} |M_{++,++}^U|^2. \quad (54)$$

This is the unnatural parity part of N_T scaled by $N_T + \varepsilon N_L$, see (42). Integrating over t one arrives at a cross section σ_U defined in analogy to σ_T in (7). Evaluating this cross section for ρ production from the amplitudes given in (3) and (48) and using the GPDs \widetilde{H} described above, we find the results shown in Fig. 12. The cross section $\sigma_U(\rho)$ is rather small but in agreement with the preliminary HERMES result [35] at $Q^2 = 2.88 \text{ GeV}^2$ within an admittedly large error. For larger energies our approach will lead to even smaller values for σ_U since the valence quark contribution disappears and, as we mentioned above, the combined gluon and sea contribution is very small (the typical size of gluon plus sea quark contribution to $\sigma_U(\rho)$ is 0.013 nb). We note that for ρ production, the H1 data [25] provide values for σ_U that are compatible with zero (e.g. at $Q^2 = 3 \text{ GeV}^2$, $\sigma_U/\sigma = 0.03 \pm 0.07$) while the ZEUS results [24] are about 1.5σ above zero (e.g. at $Q^2 \simeq 3.4 \text{ GeV}^2$, $\sigma_U/\sigma = 0.03 \pm 0.02$). Both the experimental results are in agreement with our estimates within errors. An immediate consequence of the cancellation of gluon and sea contributions to the unnatural parity amplitude is that σ_U for ϕ production is very small, in fact compatible with zero. This is in agreement with the preliminary HERMES data [37] and with the H1 data [32]. Thus, there is indication from both theory and experiment that σ_U for ρ and ϕ production is small. Its neglect in σ_T seems to be justified. A larger cross section σ_U is to be expected for ω production because the combination $e_u \widetilde{H}_{\text{val}}^u + e_d \widetilde{H}_{\text{val}}^d$ occurs (see (4)) which is larger than $e_u \widetilde{H}_{\text{val}}^u - e_d \widetilde{H}_{\text{val}}^d$ given the relative sign of H_{val}^u and H_{val}^d .

One may wonder whether or not it is possible to generate a value for σ_U as large as,

say, the face value of the preliminary HERMES result [35] by using GPDs constructed from the polarized PDFs via the double distribution ansatz. In order to examine this issue we recall that the polarized PDFs are the difference of PDFs for helicity parallel and anti-parallel to that one of the proton while the unpolarized PDFs represent their sum. Suppose the gluon and sea quark contributions still cancel and assume now that the helicity-parallel distributions dominate which, in the limit $x \rightarrow 1$, follows from QCD [44]. In this case the double distribution ansatz leads to $\widetilde{H}_{\text{val}}^a = H_{\text{val}}^a$. Admittedly this is an extreme example since in all analyses [43, 45, 46] the polarized d -quark distribution is negative. Another, more moderate example is set by the assumption $\widetilde{H}_{\text{val}}^u = H_{\text{val}}^u$ and $\widetilde{H}_{\text{val}}^d = -H_{\text{val}}^d$, implying d quarks with dominantly antiparallel helicity. The results obtained from these two scenarios are also shown in Fig. 12. Obviously, it seems difficult to obtain agreement with the HERMES result [35] with GPDs constructed from the double distribution ansatz except extreme scenarios are realized in nature.

The size of the amplitude \mathcal{M}^U can be elucidated further by considering the initial state helicity correlation A_{LL} which can be measured with a longitudinally polarized beam and target. In contrast to cross sections and SDMEs where the corrections are bilinear in the \mathcal{M}^U terms and, hence, very small, the leading term in A_{LL} is an interference between the \mathcal{M}^N and the \mathcal{M}^U amplitudes. In fact, with the help of parity conservation as well as (9) and (10), one obtains (see [1])

$$A_{LL}[ep \rightarrow eVp] = 4\sqrt{1-\varepsilon^2} \frac{\text{Re} [\mathcal{M}_{++,+}^N \mathcal{M}_{++,+}^{U*}]}{N_T + \varepsilon N_L}. \quad (55)$$

We stress that here the target polarization is specified relative to the virtual photon direction while in experiment it is usually defined with respect to the lepton beam direction. The conversion from our specification to the one used in experiments leads to a factor $\cos \theta_\gamma$ in (55) [47] where θ_γ denotes the angle describing the rotation in the lepton plane from the direction of the incoming lepton to the one of the virtual photon. This angle is given by [47]

$$\cos \theta_\gamma = \sqrt{1 - \gamma^2 \frac{1 - y - y^2 \gamma^2 / 4}{1 + \gamma^2}} \simeq 1 - \frac{1}{2} \gamma^2 (1 - y). \quad (56)$$

The two parameters appearing in (56) are $y = (W^2 + Q^2 - m^2)/(s - m^2)$, one of the conventional variables of electroproduction, and $\gamma = 2x_{\text{Bj}}m/Q$. In the kinematical situation of interest γ is very small and, hence, $\cos \theta_\gamma \simeq 1$. According to (18) A_{LL} is of order $\langle k_\perp^2 \rangle / Q^2$ and, therefore, expected to be very small. In Fig. 12 our results for ρ production are shown at $W = 5$ and 10 GeV. For the lower energy the valence quark contribution generates values for A_{LL} of about 0.1 while at 10 GeV only extremely small values are found. The valence quark contribution has nearly disappeared at that energy and, as we mentioned above, the gluon and sea quark contributions cancel each other to a large extent. For instance, at $W = 5$ GeV and $Q^2 = 3 \text{ GeV}^2$, $A_{LL}(\rho)$ changes by -0.002 if the gluon and sea quarks are neglected. Because of the very small combined gluon and sea contributions we also predict very small values of $A_{LL}(\phi)$. For instance, at $W = 5(10)$ GeV and $Q^2 = 3 \text{ GeV}^2$, $A_{LL}(\phi) = -0.002(-0.007)$.

Recently, the COMPASS collaboration has measured A_{LL} for ρ production [48]. For Q^2 less than 2 GeV² COMPASS observes very small values for A_{LL} , which are compatible with zero. Their only data point for which its Q^2 is sufficiently large for application of the handbag approach, is inconclusive because of its extremely large error; it is at variance with our results by about 1σ , see Fig. 12. The HERMES results on this observable for ρ and ϕ production [49] is in agreement with our predictions. The SMC experiment [50] observes a double spin asymmetry for ρ production at $W = 15$ GeV that is compatible with our results within large errors. In passing we remark that A_{LL} is sensitive to the relative phase δ_{NU} between the amplitudes $\mathcal{M}_{++,++}^N$ and $\mathcal{M}_{++,++}^U$. Therefore, a large value of σ_U is not necessarily in contradiction with a small value of A_{LL} provided the phase is near 90°. Nevertheless, in our approach the phase is small. For instance, at $W = 5$ GeV and $Q^2 = 3$ GeV² we find $\delta_{NU} = 3.7^\circ$, i.e. a large value of σ_U would go along with a large value of A_{LL} in our approach. For instance, a scenario with $\widetilde{H}_{\text{val}}^a = H_{\text{val}}^a$ leads to $A_{LL}(\rho) = 0.14$ for this kinematical point of reference.

Let us return to the issue of the size of the combined gluon and sea contribution to the unnatural parity amplitude. Up to now we assumed that this combined contribution is very small as follows from the double distribution ansatz using current polarized PDFs. One may wonder whether these PDFs are really correct or whether the smallness of the combined gluon and sea contribution is perhaps a special feature of our double distribution ansatz for \widetilde{H} . First we note that in all current analyses of the polarized PDFs [43, 45, 46] Δg and the polarized sea have opposite signs and are rather small in magnitude⁹. In particular a large positive Δg is in conflict with measurements of the A_{LL} asymmetry in the production of jets [51] and π^0 mesons [52] in inclusive proton-proton collisions. A negative polarized sea is for instance seen in the HERMES semi-inclusive deep-inelastic scattering data [53]. Thus, we think that the main features of the polarized PDFs are correct. A small combined gluon and sea contribution from \widetilde{H} seems to be required by the relevant data too. Leaving aside the HERMES results on σ_U and A_{LL} for ρ production which receive contributions from $\widetilde{H}_{\text{val}}$, we note that all other pertinent data are small and in most cases compatible with zero. These data are $\sigma_U(\rho)$ from H1 [25] and ZEUS [24], the same cross section for ϕ production from HERMES [37] and H1 [32] and finally the A_{LL} data from Compass [48] and SMC [50]. Thus, scenarios in which the combined gluon and sea quark contribution is large in magnitude seem to be excluded by the current data. Our double distribution ansatz for \widetilde{H}^g and $\widetilde{H}_{\text{sea}}$, as naive it may be, qualitatively reproduces the main features of the data.

7 Proton helicity flip

The analysis of observables for vector-meson electroproduction measured with a transversely polarized proton target requires the proton helicity flip amplitude. The handbag

⁹ A negative Δg is also discussed in Ref. [45].

contribution to this amplitude is given by

$$\mathcal{M}_{\mu-, \mu+}^{Ni}(V) = -\frac{e}{2} \frac{\sqrt{-t}}{2m} \sum_a e_a \mathcal{C}_V^a \int_{\bar{x}_i}^1 d\bar{x} \left[\mathcal{H}_{\mu+, \mu+}^{Vi} + \mathcal{H}_{\mu-, \mu-}^{Vi} \right] E^i(\bar{x}, \xi, t), \quad (57)$$

where t_{\min} is ignored. Our choice of the phase of this amplitude is in accord with conventions exploited in [41, 47]. In general there is also a contribution from the GPD \tilde{E} feeding the amplitude $\mathcal{M}_{+-, ++}^{Ui}$. It is expected to be small and neglected in our estimate of the size of the proton helicity flip amplitude. The evaluation of \mathcal{M}^N for proton helicity flip is analogous to that of the non-flip amplitude (3) except that the convolution is now to be performed with the GPD E instead of H . The construction of E through double distributions is also analogous to that of H , see Sect. 3. The only but crucial difference is that the forward limit $e(x) = E(x, 0, 0)$ is inaccessible in deep inelastic lepton-nucleon scattering. However, the forward limits of the valence quark GPDs have been determined phenomenologically in the form factor analysis performed in Ref. [9]. The parameters of e_{val}^u and e_{val}^d expanded according to (51), are taken from [9]. They are quoted in Tab. 4 at a scale of 4 GeV². For other scales these functions are unknown which does not matter here since we will estimate proton helicity flip only for photon virtualities near that value. Note that in contrast to u_{val} and d_{val} , e_{val}^u and e_{val}^d have opposite signs. This is due to the fact that they are normalized by

$$\int_0^1 dx e_{\text{val}}^a(x) = \kappa_a, \quad (58)$$

where κ^a gives the contribution of quark flavor a to the anomalous magnetic moment of the proton ($\kappa_u \simeq 1.67$, $\kappa_d \simeq -2.03$). The forward limits of E for gluons and sea quarks are unknown. But there is a sum rule

$$\int_0^1 dx x e^g(x) = -\sum_a \int_0^1 dx x e_{\text{val}}^a(x) - 2 \sum_a \int_0^1 dx x e^{\bar{a}}(x), \quad (59)$$

which follows from a combination of Ji's sum rule and the momentum sum rule of deep inelastic lepton-nucleon scattering [12]. Neglecting a possible difference between e^s and $e^{\bar{s}}$, we can evaluate the valence-quark term in the sum rule (59) from the GPDs specified in Tab. 4. We obtain

$$\sum_a \int_0^1 dx x e_{\text{val}}^a(x) = 0.008 \pm 0.007. \quad (60)$$

This signals a remarkable compensation between the second moments of e_{val}^u and e_{val}^d

$$\frac{\int_0^1 dx x [e_{\text{val}}^u + e_{\text{val}}^d]}{\int_0^1 dx x [e_{\text{val}}^u - e_{\text{val}}^d]} \simeq 0.026, \quad (61)$$

which is even stronger than that of their first moments ($(\kappa_u + \kappa_d)/(\kappa_u - \kappa_d) \simeq 0.1$). The error in (60) has been estimated from those quoted in [9]. Hence, the moment of e^g in (60) is only about as large as the sum of the sea quark moments. This is to be contrasted

with the situation for H where the corresponding gluon moment is more than four times larger than the sum of the sea quark ones. Another argument that points into the same direction is the behavior of the gluon (or Pomeron) Regge trajectory. As is well-known this trajectory couples mainly to the proton helicity non-flip vertex while the flip coupling is very small. It is hard to find phenomenological evidence for a non-zero flip coupling [54]. Thus, the relative importance of gluon and valence quark GPDs is very different for E and H . It seems unlikely that E^g plays an analogously prominent role as H^g . In a first step we therefore assume that, for HERMES kinematics, proton helicity flip is dominated by the valence quarks (see also the discussion in Ref. [16]). Since the same Regge poles contribute to E and H , we therefore use the standard valence-quark trajectory here as well and assume $b_{\text{val}}^e = 0$, too. With regard to this situation we cannot estimate the size of proton helicity flip for ϕ production but we expect it to be very small. We stress that due to the opposite signs of E_{val}^u and E_{val}^d , their contribution to ρ production off protons, $\propto e_u E_{\text{val}}^u - e_d E_{\text{val}}^d$, is much smaller than that from the corresponding contribution of H_{val}^a . This provides additional justification for the neglect of E in the proton helicity non-flip amplitude (see discussion after Eq. (2)). An interesting case is ω production since for it the combinations $e_u E_{\text{val}}^u + e_d E_{\text{val}}^d$ and $e_u H_{\text{val}}^u + e_d H_{\text{val}}^d$ occur. The first combination is larger, the second one smaller than for ρ production and, hence, a markedly larger ratio of proton helicity flip and non-flip is expected for ω production. For instance, at $Q^2 = 4 \text{ GeV}^2$ and $W = 5 \text{ GeV}$ the flip/non-flip ratio of the absolute values of the ω amplitudes is about 13 times larger than the corresponding ratio for ρ production.

Recently the formalism for the SDMEs in the case of a proton target polarized perpendicular ('normal') with respect to the plane in which the scattering $\gamma^* p \rightarrow V p$ takes place, has been developed [41]. These SDMEs are denoted by $n_{\mu\mu'}^{\sigma\sigma'}$ and related to bilinear combinations of the amplitudes for helicities μ, μ' and σ, σ' of the virtual photon and the meson, respectively:

$$n_{\mu\mu'}^{\sigma\sigma'} = \frac{1}{N_T + \varepsilon N_L} \sum_{\nu'} \left[\mathcal{M}_{\sigma\nu', \mu+} \mathcal{M}_{\sigma'\nu', \mu'-}^* - \mathcal{M}_{\sigma\nu', \mu-} \mathcal{M}_{\sigma'\nu', \mu'+}^* \right]. \quad (62)$$

If one neglects the amplitudes \mathcal{M}^U in accord with our findings described in Sect. 6, as well as helicity flip transitions $\gamma^* \rightarrow V$ only the SDMEs

$$n_{\mu\mu'}^{\mu\mu'} = \frac{2}{N_T + \varepsilon N_L} \sum_{\nu'} \mathcal{M}_{\mu\nu', \mu+}^N \mathcal{M}_{\mu'\nu', \mu'-}^{N*} \quad (63)$$

are non-zero. Explicitely these SDME read

$$\begin{aligned} n_{00}^{00} &= \frac{4i}{N_T + \varepsilon N_L} \text{Im} \left[\mathcal{M}_{0-,0+}^N \mathcal{M}_{0+,0+}^{N*} \right] \\ n_{++}^{++} &= n_{-+}^{-+} = \frac{4i}{N_T + \varepsilon N_L} \text{Im} \left[\mathcal{M}_{+-,++}^N \mathcal{M}_{++,++}^{N*} \right] \\ n_{0+}^{0+} &= -(n_{+0}^{+0})^* = \frac{2}{N_T + \varepsilon N_L} \left[\mathcal{M}_{0-,0+}^N \mathcal{M}_{++,++}^{N*} - \mathcal{M}_{0+,0+}^N \mathcal{M}_{+-,++}^{N*} \right]. \end{aligned} \quad (64)$$

For non-zero SDMEs $n_{\mu\mu'}^{\mu\mu'}$ phase differences between the proton helicity flip and non-flip amplitudes are mandatory. Such phase differences are provided by the handbag approach since the non-flip amplitudes are built up by gluons and quarks while the flip amplitudes receive only contributions from the valence quarks in our model for the GPD E . Indeed we obtain the values 38.8° and 34.7° for the phase between the proton flip and non-flip amplitudes in the case of longitudinal and transverse photon polarization, respectively. In Fig. 13 the SDMEs (64) are shown versus Q^2 at $W = 5$ GeV, the trivial factor $\sqrt{-t}/2m$, see (57), is pulled out and t set to zero otherwise. The scaled SDMEs are small, of the order of five percent. These SDMEs will be measured by HERMES.

One may also consider transverse proton polarization lying in the $\gamma^*p \rightarrow Vp$ plane ('sideways'). In this case SDMEs, denoted by $s_{\mu\mu'}^{\sigma\sigma'}$ [41], occur that are analogous to (62) but with a plus sign between the two terms. The SDMEs $s_{\mu\mu'}^{\mu\mu'}$ for photon-meson helicity non-flip are given by products of two small amplitudes, $\mathcal{M}_N^{\mu\mu'}$ for proton helicity flip and \mathcal{M}^U . They are therefore very small in our approach.

Finally, we estimate the asymmetry A_{UT} of $ep \rightarrow eVp$ for a transversely polarized target, normal to the $\gamma^*p \rightarrow Vp$ scattering plane. It is measured as the $\sin(\phi - \phi_S)$ moment of the electroproduction cross section where ϕ is the azimuthal angle between the lepton and hadron plane and ϕ_S the azimuthal angle of the target spin vector defined with respect to the direction of the virtual photon [41]. As for the asymmetry A_{LL} the conversion of this spin vector into the one used in the experimental setup where the target polarization is specified relative to the lepton beam, again leads to a factor $\cos\theta_\gamma$ in principle. According to the discussion following Eq. (56) it is replaced by one. In the handbag approach the dominant contribution to this asymmetry reads

$$A_{UT}(ep \rightarrow eVp) = 4 \frac{\text{Im}[\mathcal{M}_{+-,++}^N \mathcal{M}_{++,,++}^{N*}] + \varepsilon \text{Im}[\mathcal{M}_{0-,0+}^N \mathcal{M}_{0+,0+}^{N*}]}{N_T + \varepsilon N_L} \quad (65)$$

It is just the imaginary part of the sum of n_{++}^{++} and εn_{00}^{00} and is also proportional to $\sqrt{-t}/2m$. We again pull out the latter factor and display the scaled asymmetry, evaluated at $t = 0$, in Fig. 13. We obtain a positive asymmetry. In contrast to A_{LL} it is finite to leading-twist order which is obtained from (65) by neglecting the contributions from transverse photons and evaluating those from longitudinal photons in collinear approximation. For comparison the leading-twist contribution is also shown in Fig. 13. It is not too different from the full result. A preliminary HERMES result [55] for ρ production, integrated on the range $0 \geq -t \geq 0.4$ GeV², is -0.033 ± 0.058 at $Q^2 = 3.07$ GeV² and $W = 5$ GeV while we find 0.02 ± 0.01 for this kinematical situation. Note that the scaled asymmetry is still t dependent although mildly. Ignoring this and integrating just $\sqrt{-t}$ one makes an error of about 10% at $Q^2 \simeq 3 - 4$ GeV². For ω production A_{UT} is about 10 times larger than for ρ production. For ϕ production, on the other hand, we expect a very small asymmetry since the gluon and sea contributions are not only small but cancel each other to some extent, see (59).

The asymmetry A_{UL} for an unpolarized beam and a longitudinally polarized target is given by the same expression as A_{UT} . Only the mentioned conversion factor $\cos\theta_\gamma$ is to




Figure 13: Left: The imaginary parts of the SDMEs n_{00}^{00} (solid), n_{++}^{++} (dashed) and n_{0+}^{0+} (dash-dotted line), scaled by $2m/\sqrt{-t}$, for ρ production versus Q^2 at $W = 5$ GeV. The error band is only shown for n_{00}^{00} . Right: The asymmetry A_{UT} , scaled by $2m/\sqrt{-t}$, for ρ production versus Q^2 at $W = 5$ GeV. The dashed line represents the leading-twist contribution.

be replaced by $\sin\theta_\gamma$ which is very small [47]. The beam asymmetry A_{LU} obtained with a longitudinally polarized beam and an unpolarized target is zero given that helicity flip $\gamma^* \rightarrow V$ transitions can be neglected.

8 Summary

Together with Ref. [3] this work gives an exhaustive description of light vector-meson electroproduction within the handbag approach for a wide range of kinematics reaching from the HERMES up to the HERA kinematical settings. Our handbag approach includes power corrections which suppress the leading-twist amplitude for $\gamma_L^* \rightarrow V_L$ transitions and allows for a calculation of the transverse amplitude $\gamma_T^* \rightarrow V_T$. In order to specify our handbag approach fully we have to mention the soft physics input, namely the GPDs that are constructed from PDFs with the help of double distributions, and the light-cone wavefunctions for the mesons. GPDs and wavefunctions affect the handbag amplitudes differently and can therefore be disentangled. The wavefunctions provide effects of order $\langle \mathbf{k}_\perp^2 \rangle / Q^2$ controlled by the transverse size parameter a_V while the GPDs mainly influence the x_{Bj} dependence of the amplitudes or, at fixed Q^2 , the W dependence. Besides the dominant contributions from the GPD H ('natural parity' contribution) we also estimated effects from the GPDs \widetilde{H} ('unnatural parity' contribution) and E controlling the proton helicity flip amplitudes. These effects are generally small. With our analysis we achieve a very good description of the HERA, HERMES and COMPASS data on the separated and unseparated cross sections for ρ and ϕ electroproduction, on the ratio σ_L/σ_T , on the SDMEs and on some spin asymmetries. The only problem we observed is that the relative

phase between the longitudinal and transverse amplitudes seems to be larger in experiment, in particular in the HERMES experiment [35], than our handbag approach predicts. The neglect of contributions from transitions other than $\gamma_L^* \rightarrow V_L$ and $\gamma_T^* \rightarrow V_T$ to the cross sections seems to be justified. Only little contributions from $\gamma_T^* \rightarrow V_L$ transitions are to be observed in some of the SDMEs experimentally. We note that in Ref. [56] the longitudinal amplitude has also been analyzed within the handbag approach. The main difference to our work is that in [56] the gluonic contribution is treated in leading-log approximation [57] and added incoherently to the quark amplitudes. This line of action underestimates the gluonic contribution at low energies.

The applicability of our approach is limited to $Q^2 \gtrsim 3 - 4 \text{ GeV}^2$, $W \gtrsim 4 - 5 \text{ GeV}$ and $x_{\text{Bj}} \lesssim 0.2$. The restriction of Q^2 is due to the mentioned - and still unsettled - difficulties with higher-order perturbative corrections as well as due to the neglected corrections of order m^2/Q^2 and $-t/Q^2$. There may be also power corrections of other dynamical origin which become large at low Q^2 . The restriction of W has its origin in the asymmetric minimum the cross sections exhibit at $W \simeq 3 - 4 \text{ GeV}$. While the cross sections [24, 25, 34, 58] mildly increase towards larger W , they [35, 59, 60] increase sharply in the opposite direction. In fact they rise by nearly an order of magnitude between $W \simeq 4$ and 2 GeV . Obviously, a new dynamical mechanism seems to set in and the handbag physics is perhaps not applicable here. On the other side, it dominates for $W \gtrsim 4 \text{ GeV}$. The mild increase of the cross sections with energy beyond the minimum is well described by the handbag physics as the results presented in this article and in Ref. [3] reveal. The restriction of x_{Bj} is of difference quality. It allows to neglect contributions of order x_{Bj}^2 or ξ^2 , e.g. in (1), which simplifies the analysis of vector-meson electroproduction strongly.

Acknowledgements

We thank A. Borissov, M. Diehl, N. d’Hose, A. Levy, W.-D. Nowak and A. Sandacz for discussions. We are also grateful to the HERMES and COMPASS collaborations for permission to use preliminary data. This work has been supported in part by the Russian Foundation for Basic Research, Grant 06-02-16215, the Integrated Infrastructure Initiative “Hadron Physics” of the European Union, contract No. 506078 and by the Heisenberg-Landau program.

References

- [1] S. V. Goloskokov and P. Kroll, Eur. Phys. J. C **42**, 281 (2005) [hep-ph/0501242].
- [2] J. Botts and G. Sterman, Nucl. Phys. B **325**, 62 (1989).
- [3] S. V. Goloskokov and P. Kroll, Eur. Phys. J. C **50**, 829 (2007) [hep-ph/0611290].
- [4] L. Mankiewicz and G. Piller, Phys. Rev. D **61**, 074013 (2000) [hep-ph/9905287].

- [5] I. V. Anikin and O. V. Teryaev, Phys. Lett. B **554**, 51 (2003) [hep-ph/0211028].
- [6] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. Nadolsky and W. K. Tung, JHEP **0207**, 012 (2002) [hep-ph/0201195].
- [7] D. Mueller *et al.*, Fortsch. Phys. **42**, 101 (1994) [hep-ph/9812448].
- [8] A. V. Radyushkin, Phys. Lett. B **449**, 81 (1999) [hep-ph/9810466].
- [9] M. Diehl, T. Feldmann, R. Jakob and P. Kroll, Eur. Phys. J. C **39**, 1 (2005) [hep-ph/0408173].
- [10] A.V. Radyushkin, Phys. Lett. B **385**, 333 (1996) [hep-ph/9605431].
- [11] J.C. Collins, L. Frankfurt and M. Strikman, Phys. Rev. D **56**, 2982 (1997) [hep-ph/9611433].
- [12] M. Diehl, Phys. Rept. **388**, 41 (2003) [hep-ph/0307382].
- [13] M. Diehl, W. Kugler, A. Schafer and C. Weiss, Phys. Rev. D **72**, 034034 (2005)[hep-ph/0506171].
- [14] M. Diehl, W. Kugler, A. Schafer and C. Weiss, Phys. Rev. D **72**, 059902 (2005) [Erratum].
- [15] D. Yu. Ivanov, L. Szymanowski and G. Krasnikov, JETP Lett. **80**, 226 (2004) [Pisma Zh. Eksp. Teor. Fiz. **80**, 255 (2004)] [hep-ph/0407207].
- [16] M. Diehl and W. Kugler, arXiv:0708.1121 [hep-ph].
- [17] D. Yu. Ivanov, to be published in the proceedings of the EDS07 conference, Hamburg (2007).
- [18] S. Catani and F. Hautmann, Nucl. Phys. B **427**, 475 (1994) [hep-ph/9405388].
- [19] R. Jakob and P. Kroll, Phys. Lett. B **315**, 463 (1993) [hep-ph/9306259].
- [20] R. Jakob and P. Kroll, Phys. Lett. B **319**, 545 (1993) [Erratum].
- [21] P.V. Landshoff, J.C. Polkinghorne and R.D. Short, Nucl. Phys. B **28**, 225 (1971).
- [22] R. P. Feynman, Photon-Hadron Interactions (W.A. Benjamin, INC., Reading, Massachusetts, 1972).
- [23] M. V. Polyakov and C. Weiss, Phys. Rev. D **60**, 114017 (1999) [hep-ph/9902451].
- [24] S. Chekanov [ZEUS Collaboration], arXiv:0708.1478 [hep-ex].
- [25] C. Adloff *et al.* [H1 Collaboration], Eur. Phys. J. **C13**, 371 (2000) [hep-ex/9902019].

- [26] S. Chekanov *et al.* [ZEUS Collaboration], Nucl. Phys. B **718**, 3 (2005) [hep-ex/0504010].
- [27] A. V. Vinnikov, hep-ph/0604248.
- [28] P. Ball and V. M. Braun, Phys. Rev. D **54**, 2182 (1996) [hep-ph/9602323].
- [29] M. A. Shifman and M. I. Vysotsky, Nucl. Phys. B **186**, 475 (1981).
- [30] M. Glück, C. Pisano and E. Reya, Eur. Phys. J. C **50**, 29 (2007) [hep-ph/0610060].
- [31] M. Glück, E. Reya and A. Vogt, Eur. Phys. J. C **5**, 461 (1998) [hep-ph/9806404].
- [32] C. Adloff *et al.* [H1 Collaboration], Phys. Lett. B **483**, 360 (2000) [hep-ex/0005010].
- [33] S. Aid *et al.* [H1 Collaboration], Nucl. Phys. B **468**, 3 (1996) [hep-ex/9602007].
- [34] J. Breitweg *et al.* [ZEUS Collaboration], Eur. Phys. J. C **6**, 603 (1999) [hep-ex/9808020].
- [35] A. Borissov [HERMES Collaboration], Proceedings of "Diffraction 06", PoS (DIFF2006) 014.
- [36] B. Marianski [HERMES collaboration], Proceedings of "DIS 2006", World Scientific, 2007.
- [37] A. Borissov [HERMES collaboration], Proceedings of DSPIN-07, to be published.
- [38] D. Neyret [COMPASS collaboration] preliminary data presented at SPIN2004, Trieste
- [39] J. Breitweg *et al.*, [ZEUS Collaboration], Eur. Phys. J. C **12**, 393 (2000) [hep-ex/9908026].
- [40] K. Schilling and G. Wolf, Nucl. Phys. B **61**, 381 (1973).
- [41] M. Diehl, JHEP0709, 064 (2007) [arXiv:0704.1565].
- [42] M. Diehl and D. Y. Ivanov, arXiv:0707.0351 [hep-ph].
- [43] J. Blümlein and H. Böttcher, Nucl. Phys. B **636**, 225 (2002) [hep-ph/0203155].
- [44] S. J. Brodsky, M. Burkardt and I. Schmidt, Nucl. Phys. B **441**, 197 (1995) [hep-ph/9401328].
- [45] E. Leader, A. V. Sidorov and D. B. Stamenov, Phys. Rev. D **75**, 074027 (2007) [hep-ph/0612360].
- [46] M. Hirai, S. Kumano and N. Saito, Phys. Rev. D **74**, 014015 (2006) [hep-ph/0603213].
- [47] M. Diehl and S. Sapeta, Eur. Phys. J. C **41**, 515 (2005) [hep-ph/0503023].

- [48] V. Y. Alexakhin *et al.* [COMPASS Collaboration], Eur. Phys. J. C **52**, 255 (2007) [arXiv:0704.1863].
- [49] A. Airapetian *et al.* [HERMES Collaboration], Eur. Phys. J. C **29**, 171 (2003) [hep-ex/0302012].
- [50] A. Tripet [Spin muon collaboration], Nucl. Phys. Proc. Suppl. **79**, 529 (1999) [hep-ex/9906008].
- [51] K. Kowalik [STAR Collaboration], arXiv:0706.2667 [hep-ex].
- [52] A. Adare *et al.* [PHENIX Collaboration] Phys. Rev. D **76**, 051106 (2007) [arXiv:0704.3599].
- [53] A. Airapetian *et al.* [HERMES Collaboration], Phys. Rev. D **71**, 012003 (2005) [hep-ex/0407032].
- [54] A. Donnachie Phys. Lett. B **611**, 255 (2005) [hep-ph/0412085].
- [55] A. Rostomyan [HERMES Collab.], Proceedings of "DIS 2007", World Scientific, 2007.
- [56] M. Vanderhaeghen, P. A. M. Guichon and M. Guidal, Phys. Rev. D **60**, 094017 (1999) [hep-ph/9905372].
- [57] L. Frankfurt, W. Koepf and M. Strikman, Phys. Rev. D **54**, 3194 (1996) [arXiv:hep-ph/9509311].
- [58] M. R. Adams *et al.* [E665 Collaboration], Z. Phys. C **74**, 237 (1997).
- [59] D. G. Cassel *et al.*, Phys. Rev. D **24**, 2787 (1981).
- [60] C. Hadjidakis *et al.* [CLAS Collaboration], Phys. Lett. B **605**, 256 (2005) [hep-ex/0408005].